


AHPWA1DUSA

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1 (Currently Amended). A method of hormone replacement therapy, the method comprising administering to a mammal in need thereof a compound of the formula (I):



wherein:

R¹ and R² are independent substituents selected from the group consisting of H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₂ to C₆ alkenyl, substituted C₂ to C₆ alkenyl, C₂ to C₆ alkynyl, substituted C₂ to C₆ alkynyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, COR^A, and NR^BCOR^A;

or R¹ and R² are fused to form:

- a) a carbon-based 3 to 8 membered saturated spirocyclic ring;
- b) a carbon-based 3 to 8 membered spirocyclic ring having in its backbone one or more carbon-carbon double bonds; or
- c) a carbon-based 3 to 8 membered heterocyclic ring having in its backbone one to three heteroatoms selected from the group consisting of O, S and N;

AHPWA1DUSA

the spirocyclic rings of a), b) and c) being optionally substituted by from 1 to 4 groups selected from the group consisting of fluorine, C₁ to C₆ alkyl, C₁ to C₆ alkoxy, C₁ to C₆ thioalkyl, CF₃, OH, CN, NH₂, NH(C₁ to C₆ alkyl), and N(C₁ to C₆ alkyl)₂;

R^A is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^B is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;


R³ is H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₆ alkenyl, substituted C₃ to C₆ alkenyl, alkynyl, substituted alkynyl, or COR^C;

R^C is H, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, aryl, substituted aryl, C₁ to C₄ alkoxy, substituted C₁ to C₄ alkoxy, C₁ to C₄ aminoalkyl, or substituted C₁ to C₄ aminoalkyl;

R⁴ is H, halogen, CN, NO₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, alkynyl, substituted alkynyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, amino, C₁ to C₆ aminoalkyl, or substituted C₁ to C₆ aminoalkyl;

R⁵ is selected from the group consisting of (i) and (ii):

- (i) a substituted benzene ring having the substituents X, Y and Z as shown below:



wherein:

X is selected from the group consisting of H, halogen, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkoxy, substituted C₁ to C₃ thioalkoxy, amino, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 or 6 membered heterocyclic ring containing in its backbone 1 to 3 heteroatoms selected from the group consisting of O, S, and N, COR^D, OCOR^D, and NR^ECOR^D;

AHPWA1DUSA

R^D is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

Y and Z are independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, aminoalkyl, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkoxy;

wherein X, Y, and Z are not all H; and

(ii) a five or six membered ring having in its backbone 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and containing one or two independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, C₁ to C₄ alkyl, C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, COR^F, and NR^GCOR^F;

R^F is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^G is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R^6 is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

wherein when R⁵ is a five-membered ring having in its backbone a NR⁶ heteroatom, and when R⁵ is attached at the two position of said ring, there is no CN substituent in the five position on said ring;

or pharmaceutically acceptable salt thereof.

2(Original). The method according to Claim 1, wherein:

R^1 is H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, COR^A, or NR^BCOR^A;

R^2 is H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₂ to C₆ alkenyl, substituted C₂ to C₆ alkenyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, COR^A, or NR^BCOR^A;

AHPWA1DUSA

R⁵ is (i) or (ii):

(i) the substituted benzene ring, wherein:

X is selected from the group consisting of halogen, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkoxy, substituted C₁ to C₃ thioalkoxy, amino, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing in its backbone 1 to 3 heteroatoms selected from the group consisting of N, O, and S, COR^D, OCOR^D, and NR^ECOR^D;

Y and Z are independent substituents selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, and C₁ to C₃ thioalkoxy; or

(ii) the five or six membered ring, wherein said one or two independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

R⁶ is H or C₁ to C₃ alkyl.

3(Original). The method according to Claim 1, wherein:

R¹ is H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, COR^A, or NR^BCOR^A;

R⁴ is H, halogen, CN, NO₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, amino, C₁ to C₆ aminoalkyl, or substituted C₁ to C₆ aminoalkyl;

R⁵ is (iii) or (iv):

(iii) the substituted benzene ring, wherein

X is selected from the group consisting of halogen, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkoxy, substituted C₁ to C₃ thioalkoxy, amino, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing in

AHPWA1DUSA

its backbone 1 to 3 heteroatoms selected from the group consisting of N, O, and S, COR^D, OCOR^D, and NR^ECOR^D;

Y and Z are independent substituents selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, and C₁ to C₃ thioalkoxy; or

(iv) the five or six membered ring, wherein said ring contains one or two independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

R⁶ is H or C₁ to C₃ alkyl.

4(Original). The method according to Claim 1, wherein:

R¹ = R² and are selected from the group consisting of C₁ to C₃ alkyl and substituted C₁ to C₃ alkyl, or R¹ and R² are fused to form the carbon-based 3 to 6 membered saturated spirocyclic ring;


R³ is H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, or COR^C;

R^C is H, C₁ to C₄ alkyl, or C₁ to C₄ alkoxy;

R⁴ is H, halogen, CN, NO₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, or substituted C₁ to C₃ alkoxy;

R⁵ is (v), (vi), or (vii):

(v) the substituted benzene ring of the structure:




wherein:

X is halogen, CN, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing in its backbone 1 to 3 heteroatoms selected from the group consisting of N, O, and S, or C₁ to C₃ thioalkoxy;

Y is H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkoxy;

AHPWA1DUSA


(vi) the five membered ring having the structure:




wherein:

U is O, S, or NR⁶;X' is selected from the group consisting of halogen, CN, NO₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;Y' is selected from the group consisting of H and C₁ to C₄ alkyl; or


(vii) the six membered ring having the structure:



wherein:

X¹ is N or CX²;X² is halogen, CN, C₁ to C₃ alkoxy, or NO₂.5(Original). The method according to Claim 4, wherein R⁵ is selected from the group consisting of:

AHPWA1DUSA




6(Original). The method according to Claim 1, wherein:
 R^1 and R^2 are CH_3 or R^1 and R^2 are fused to form the carbon-based 6 membered saturated spirocyclic ring;

R^3 is H, OH, NH_2 , CH_3 , substituted CH_3 , or COR^C ;

R^C is H, C_1 to C_3 alkyl, or C_1 to C_4 alkoxy;

R^4 is H, halogen, NO_2 , CN, or C_1 to C_3 alkyl;

R^5 is the substituted benzene ring having the formula:



wherein:

X is selected from the group consisting of halogen, CN, methoxy, NO_2 , and the five-membered heterocyclic ring, wherein said ring is 2-thiazole;

Y is H or halogen, wherein said halogen is F.

7(Original). The method according to Claim 1, wherein:
 R^1 and R^2 are CH_3 or R^1 and R^2 are fused to form the carbon-based 6 membered saturated spirocyclic ring;


R^3 is H, OH, NH_2 , CH_3 , substituted CH_3 , or COR^C ;

R^C is H, C_1 to C_3 alkyl, or C_1 to C_4 alkoxy;

R^4 is H, halogen, NO_2 , CN, or C_1 to C_3 alkyl;

R^5 is the five membered ring having the structure:

AHPWA1DUSA



wherein:

U is O, S, or NH;

X' is halogen, CN, or NO₂, provided that when **U** is NH, then **X'** is not CN;

Y' is H or C₁ to C₄ alkyl.

8(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 6-(3-Chlorophenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one;
- b) 6-(3-Methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
- c) 6-(2-Chloro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- d) 6-(4-Chloro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
- e) 6-(3-Chloro-phenyl)-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- f) 6-(3-Chloro-phenyl)-4-ethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; and
- g) 6-(3-Chloro-phenyl)-4-phenyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
or a pharmaceutically acceptable salt thereof.

9(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile;
- b) 4,4-Dimethyl-6-(3-nitrophenyl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one;
- c) 6-(3-Bromo-5-fluorophenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one;

AHPWA1DUSA

- d) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluorobenzonitrile;
- e) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-nicotinonitrile;
- f) 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile;
- g) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile;
- h) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-methyl-thiophene-2-carbonitrile;
- i) 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbonitrile; and
- j) 4,4-Diethyl-6-(3-nitrophenyl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one; or a pharmaceutically acceptable salt thereof.

10(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 6-(3-Chlorophenyl)-4,4-diethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one;
- b) 6-(3-Chlorophenyl)-spiro[4H-3,1-benzoxazine-4,1'-cyclohexane]-2-(1H)-one;
- c) 6-(3-Chlorophenyl)-spiro-[4H-3,1-benzoxazine-4,1'-cyclopentane]-2(1H)-one;
- d) 6-(3-Nitrophenyl)-spiro[4H-3,1-benzoxazine-4,1'-cyclohexan]-2(1H)-one;
- e) 4-Allyl-6-(3-chlorophenyl)-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- f) 6-(3-Chlorophenyl)-4-methyl-4-propyn-1-yl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; and

AHPWA1DUSA

g) 6-(3-Chlorophenyl)-4-ethynyl-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
or a pharmaceutically acceptable salt thereof.

11(Original). The method according to Claim 1, wherein said compound is selected from the group of:

- a) 6-(3-Chlorophenyl)-4-methyl-4-phenyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - b) 4-Benzyl-6-(3-chloro-phenyl)-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - c) 6-(3-Chloro-phenyl)-4-cyclopropyl-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - d) 6-(3-Chloro-phenyl)-4-cyclopropyl-4-propyn-1-yl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - e) 6-(3-Chloro-phenyl)-4,4-dicyclopropyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - f) 6-(3-Chloro-phenyl)-4,4-dipropyn-1-yl-1,4-dihydrobenzo[d][1,3]oxazin-2-one;
 - g) 6-(3-Bromo-5-fluorophenyl)-1,4,4-trimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; and
 - h) 6-(3-Methoxyphenyl)-4-methyl-4-trifluoromethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- or a pharmaceutically acceptable salt thereof.

12(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 6-(3-Acetyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;

AHPWA1DUSA

- b) 6-(3-Benzoyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
- c) 4-(4,4-Dicyclopropyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile;
- d) 6-(3-Bromo-5-fluoro-phenyl)-4,4-dicyclopropyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one;
- e) 3-(4,4-Dicyclopropyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile;
- f) 6-(3-Bromo-5-methyl-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one;
- g) 6-(3-Bromo-5-trifluoromethoxy-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; and
- h) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-methyl-benzonitrile;
or a pharmaceutically acceptable salt thereof.

13(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-trifluoromethoxy-benzonitrile;
- b) 6-(3,5-difluoro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one;
- c) 6-(3,5-dichloro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one;
- d) 6-(3,5-Bis-trifluoromethyl-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one;
- e) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-methoxy-benzonitrile;

AHPWA1DUSA

- f) 6-(3-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
 - g) 6-(3-Chloro-4-fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - h) 3-(1-Diethoxymethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile; and
 - i) 3-Fluoro-5-(1-methoxymethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile;
- or a pharmaceutically acceptable salt thereof.

14(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) Phosphoric acid 6-(3-cyano-5-fluoro-phenyl)-4,4-dimethyl-4H-benzo[d][1,3]oxazin-2-yl ester diethyl ether;
 - b) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-fluoro-benzonitrile;
 - c) 6-(3-Chloro-4-fluoro-phenyl)-8-fluoro-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one;
 - d) 6-(3-Bromo-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
 - e) 6-(3-Ethynyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
 - f) 3-[3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-phenyl]-propynenitrile;
 - g) 6-(3-Fluoro-5-nitro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; and
 - h) 6-(3-Chloro-5-fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- or a pharmaceutically acceptable salt thereof.

AHPWA1DUSA

15(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 3-Chloro-5-(4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile;
 - b) 6-(3,5-Dinitro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - c) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-isophthalonitrile;
 - d) 4,4-Dimethyl-6-(3-thiazol-2-yl-phenyl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - e) 6-(3-Fluoro-5-methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - f) 6-(3-Fluoro-5-trifluoromethyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - g) 6-(5-Bromo-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - h) 6-(3-Cyano-5-fluoro-phenyl)-4,4-dimethyl-2-oxo-4H-benzo[d][1,3]oxazine-1-carboxylic acid tert-butyl ester; and
 - i) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluoro-benzonitrile;
- or a pharmaceutically acceptable salt thereof.

16(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 4-(8-Fluoro-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile;
- b) 3-Fluoro-5-(8-fluoro-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile;

AHPWA1DUSA

- c) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-3-carbonitrile;
- d) 2-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-3-carbonitrile;
- e) 6-(1,2,4-thiadiazol-3-yl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- f) 6-(3-Fluoro-5-thiophen-3-yl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- g) 2-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-pyrrole-1-carboxylic acid tert-butyl ester;
- h) 2-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-nitro-pyrrole-1-carboxylic acid tert-butyl ester;
- i) 4,4-Dimethyl-6-(5-nitro-1H-pyrrol-2-yl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one; and
- j) 4,4-Dimethyl-6-(1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
or a pharmaceutically acceptable salt thereof.

17(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 4,4-Dimethyl-6-(1-methyl-1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- b) 4,4-Dimethyl-6-(1-methyl-5-nitro-1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- c) 3-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-benzonitrile;
- d) 3-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-5-fluorobenzonitrile;

AHPWA1DUSA

- e) 4-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-2-thiophenecarbonitrile;
- f) 5-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-2-thiophenecarbonitrile;
- g) 5-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-4-methyl-2-thiophenecarbonitrile;
- h) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-ethyl-thiophene-2-carbonitrile; and
- i) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-n-propyl-thiophene-2-carbonitrile;
or a pharmaceutically acceptable salt thereof.

18(Original). The method according to Claim 1, wherein said compound is selected from the group consisting of:

- a) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-n-butyl-thiophene-2-carbonitrile;
- b) 6-(4-Cyano-3-fluoro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one;
- c) 6-(4-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
- d) 6-(3,4-Difluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- e) 6-(2-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one;
- f) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile;
- g) 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbonitrile; and

AHPWA1DUSA

- h) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluoro-benzonitrile;
or a pharmaceutically acceptable salt thereof.

19(Original). The method according to claim 1, wherein said compound is selected from the group consisting of:

- a) 6-(3-Methoxyphenyl)spiro[4H-3,1-benzoxazine-4,1-cyclobutan]-2(1H)-one;
- b) 8-Bromo-6-(3-chloro-4-fluorophenyl)-4,4-dimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one;
- c) 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile;
- d) 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile;
- e) 5-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-2-fluorobenzonitrile;
- f) 6-(3-Bromophenyl)-1,4,4-trimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one;
- g) 6-(3-Fluorophenyl)-4-methyl-1,4-dihydro-2H-3,1-benzoxazin-2-one;
- h) 3-(4,4-Dimethyl-8-methoxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile;
- i) 3-(4,4-Dimethyl-8-hydroxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile;
- j) 6-(2,3-Difluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
- k) 3-(1-Ethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile;
- l) [6-(4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)pyridin-2-yl]acetonitrile;

AHPWA1DUSA

- m) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-5-fluoro-phenylacetonitrile;
 - n) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-4-fluoro-phenylacetonitrile;
 - o) 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluorophenylacetonitrile;
 - p) 2-(4,4-Dimethyl 2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile;
 - q) 6-(3-Fluoro-4-methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one;
 - r) 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile;
 - s) 6-(6-Amino-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazine-2-one;
 - t) 6-(5-Diethoxymethyl-furan-3-yl) -4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; and
 - u) 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbaldehyde;
- or a pharmaceutically acceptable salt thereof.

20(Original). The method according to claim 1, wherein R⁵ is said five-membered ring b).

21(Original). The method according to claim 20, wherein said five-membered ring b) is a thiophene group.

22(Original). The method according to claim 20, wherein said five-membered ring b) is a furan group.

AHPWA1DUSA

23(Original). The method according to claim 20, wherein said five-membered ring b) is a pyrrole group.

24(Original). The method according to claim 20, wherein said five-membered ring b) is a thiazole group.

25(Original). The method according to claim 20, wherein said five-membered ring b) is an oxazole group.

26(Original). The method according to claim 20, wherein said five-membered ring b) is an imidazole group.

27(Original). The method according to claim 1, wherein R⁵ is said six-membered ring b).

28(Original). The method according to claim 27, wherein said six-membered ring b) is a pyridine group.

29(Original). The method according to claim 1, wherein R⁵ is said substituted benzene ring a).

30(Original). The method according to claim 29, wherein said substituted benzene ring a) is an optionally substituted phenyl group.

31(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-chloro-4-fluoro-phenyl group.

32(Original). The method according to claim 30, wherein said substituted phenyl group is a 3,5-dichloro-phenyl group.

AHPWA1DUSA

33(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-cyano-4-fluoro-phenyl group.

34(Original). The method according to claim 30, wherein said substituted phenyl group is a 3,4-difluoro-phenyl group.

35(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-cyano-5-chloro-phenyl group.

36(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-trifluoromethyl-5-fluoro-phenyl group.

37(Original). The method according to claim 30, wherein said substituted phenyl group is a 2-fluoro-3-cyano-phenyl group.

38(Original). The method according to claim 30, wherein said substituted phenyl group is a 2-fluoro-phenyl group.

39(Original). The method according to claim 30, wherein said substituted phenyl group is a 4-cyano-3-furanyl-phenyl group.

40(Original). The method according to claim 30, wherein said substituted phenyl group is a 3,4-dichloro-phenyl group.

41(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-fluoro-4-chloro-phenyl group.

42(Original). The method according to claim 30, wherein said substituted phenyl group is a 3-bromo-4-fluoro-phenyl group.


AHPWA1DUSA

43(Original). The method according to claim 30, wherein said substituted phenyl group is a 3,5-dibromo-phenyl group.

44(Original). The method according to claim 1, wherein R¹ and R² are C₁ to C₆ alkyl.

45(Original). A method of hormone replacement therapy, the method comprising administering to a mammal in need thereof a compound selected from the group consisting of 4,4-Dimethyl-6-[3-(1H-tetrazol-5-yl)-phenyl]-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 4,4-Dimethyl-6-(3-trimethylsilanylethynyl-phenyl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(5-Bromo-1-oxy-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; N-[4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-2-fluoro-phenyl]-acetamide; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl) benzenesulfonamide; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-thiophene-2-sulfonamide; and 4-(1,4-Dihydro-4,4-dimethyl-2-oxo-2H-3,1-benzoxazin-6-yl)-2-furancarboxaldehyde oxime; or pharmaceutically acceptable salt thereof.

46(Currently Amended). A method of hormone replacement therapy, the method comprising administering to a mammal in need thereof a compound of the formula (I):



I

wherein:

AHPWA1DUSA

R^1 and R^2 are independent substituents selected from the group consisting of H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₂ to C₆ alkenyl, substituted C₂ to C₆ alkenyl, C₂ to C₆ alkynyl, substituted C₂ to C₆ alkynyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, COR^A, and NR^BCOR^A;

or R^1 and R^2 are fused to form:

- a) a carbon-based 3 to 8 membered saturated spirocyclic ring;
- b) a carbon-based 3 to 8 membered spirocyclic ring having in its backbone one or more carbon-carbon double bonds; or
- c) a carbon-based 3 to 8 membered heterocyclic ring having in its backbone one to three heteroatoms selected from the group consisting of O, S and N;

the spirocyclic rings of a), b) and c) being optionally substituted by from 1 to 4 groups selected from the group consisting of fluorine, C₁ to C₆ alkyl, C₁ to C₆ alkoxy, C₁ to C₆ thioalkyl, CF₃, OH, CN, NH₂, NH(C₁ to C₆ alkyl), and N(C₁ to C₆ alkyl)₂;

R^A is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^B is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R^3 is H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₆ alkenyl, substituted C₃ to C₆ alkenyl, alkynyl, substituted alkynyl, or COR^C;


R^C is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^4 is H, halogen, CN, NO₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, alkynyl, substituted alkynyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, amino, C₁ to C₆ aminoalkyl, or substituted C₁ to C₆ aminoalkyl;

R^5 is selected from the group consisting of (i) and (ii):

- (i) a substituted benzene ring having the substituents X, Y and Z as shown below:

AHPWA1DUSA



wherein:

X is selected from the group consisting of H, halogen, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkoxy, substituted C₁ to C₃ thioalkoxy, amino, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 or 6 membered heterocyclic ring containing in its backbone 1 to 3 heteroatoms selected from the group consisting of O, S, and N, COR^D, OCOR^D, and NR^ECOR^D;

R^D is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

Y and Z are independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, aminoalkyl, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, and C₁ to C₃ thioalkoxy;

wherein X, Y, and Z are not all H; and

(ii) a five or six membered ring having in its backbone 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR^G and containing one or two independent substituents selected from the group consisting of H, halogen, CN, NO₂, amino, C₁ to C₃ alkyl, C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, COR^F, and NR^GCOR^F;

R^F is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^G is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R^G is H or C₁ to C₃ alkyl;

AHPWA1DUSA

wherein when R⁵ is a five-membered ring having in its backbone a NR⁶ heteroatom, and when R⁵ is attached at the two position of said ring, there is no CN substituent in the five position on said ring;
or a pharmaceutically acceptable salt thereof.

47(Original). The method according to Claim 46, wherein said compound is selected from the group consisting of 6-(3-Chlorophenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Methoxy-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; 6-(2-Chloro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(4-Chloro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; 6-(3-Chloro-phenyl)-4-methyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4-ethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4-phenyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile; 4,4-Dimethyl-6-(3-nitrophenyl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Bromo-5-fluorophenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluorobenzonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-nicotinonitrile; 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-methyl-thiophene-2-carbonitrile; 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbonitrile; 4,4-Diethyl-6-(3-nitrophenyl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Chlorophenyl)-4,4-diethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Chlorophenyl)-spiro[4H-3,1-benzoxazine-4,1'-cyclohexane]-2-(1H)-one; 6-(3-Chlorophenyl)-spiro-[4H-3,1-benzoxazine-4,1'-cyclopentane]-2(1H)-one; 6-(3-Nitrophenyl)-spiro[4H-3,1-benzoxazine-4,1'-cyclohexan]-2(1H)-one; 4-Allyl-6-(3-chlorophenyl)-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chlorophenyl)-4-methyl-4-propyn-1-yl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chlorophenyl)-4-

AHPWA1DUSA

ethynyl-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chlorophenyl)-4-methyl-4-phenyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 4-Benzyl-6-(3-chloro-phenyl)-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4-cyclopropyl-4-methyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4-cyclopropyl-4-propyn-1-yl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4,4-dicyclopethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chloro-phenyl)-4,4-dipropyn-1-yl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Bromo-5-fluorophenyl)-1,4,4-trimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 6-(3-Methoxyphenyl)-4-methyl-4-trifluoromethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Acetyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 6-(3-Benzoyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 4-(4,4-Dicyclopethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile; 6-(3-Bromo-5-fluoro-phenyl)-4,4-dicyclopethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one; 3-(4,4-Dicyclopethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile; 6-(3-Bromo-5-methyl-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one; 6-(3-Bromo-5-trifluoromethoxy-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-methyl-benzonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-trifluoromethoxy-benzonitrile; 6-(3,5-difluoro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one; 6-(3,5-dichloro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo-[d][1,3]oxazin-2-one; 6-(3,5-Bis-trifluoromethyl-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-methoxy-benzonitrile; 6-(3-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 6-(3-Chloro-4-fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(1-Diethoxymethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile; 3-Fluoro-5-(1-methoxymethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile; Phosphoric acid 6-(3-cyano-5-fluoro-phenyl)-4,4-dimethyl-4H-benzo[d][1,3]oxazin-2-yl ester diethyl ether; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-fluoro-benzonitrile; 6-(3-Chloro-4-fluoro-

AHPWA1DUSA

phenyl)-8-fluoro-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; 6-(3-Bromo-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 6-(3-Ethynyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 3-[3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-phenyl]-propynenitrile; 6-(3-Fluoro-5-nitro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Chloro-5-fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-Chloro-5-(4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile; 6-(3,5-Dinitro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-isophthalonitrile; 4,4-Dimethyl-6-(3-thiazol-2-yl-phenyl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Fluoro-5-methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Fluoro-5-trifluoromethyl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(5-Bromo-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(5-Bromo-1-oxy-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Cyano-5-fluoro-phenyl)-4,4-dimethyl-2-oxo-4H-benzo[d][1,3]oxazine-1-carboxylic acid tert-butyl ester; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluoro-benzonitrile; 4-(8-Fluoro-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-2-carbonitrile; 3-Fluoro-5-(8-fluoro-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-benzonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-3-carbonitrile; 2-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-thiophene-3-carbonitrile; 6-(1,2,4-thiadiazol-3-yl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(3-Fluoro-5-thiophen-3-yl-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 4,4-Dimethyl-6-(5-nitro-1H-pyrrol-2-yl)-1,4-dihydrobenzo[d][1,3]oxazin-2-one; 4,4-Dimethyl-6-(1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 4,4-Dimethyl-6-(1-methyl-1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 4,4-Dimethyl-6-(1-methyl-5-nitro-1H-pyrrol-2-yl)-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-benzonitrile; 3-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-5-fluorobenzonitrile; 4-(1,2-Dihydro-2-oxospiro[4H-


AHPWA1DUSA

3,1-benzoxazine-4,1-cyclohexan]-6-yl)-2-thiophenecarbonitrile; 5-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-2-thiophenecarbonitrile; 5-(1,2-Dihydro-2-oxospiro[4H-3,1-benzoxazine-4,1-cyclohexan]-6-yl)-4-methyl-2-thiophenecarbonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-ethyl-thiophene-2-carbonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-n-propyl-thiophene-2-carbonitrile; 6-(4-Cyano-3-fluoro-phenyl)-4,4-dimethyl-1,4-dihydrobenzo[d][1,3]-oxazin-2-one; 6-(4-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 6-(3,4-Difluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 6-(2-Fluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]-oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluoro-benzonitrile; 6-(3-Methoxyphenyl)spiro[4H-3,1-benzoxazine-4,1-cyclobutan]-2(1H)-one; 8-Bromo-6-(3-chloro-4-fluorophenyl)-4,4-dimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 5-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-2-fluorobenzonitrile; 6-(3-Bromophenyl)-1,4,4-trimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 6-(3-Fluorophenyl)-4-methyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 3-(4,4-Dimethyl-8-methoxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 3-(4,4-Dimethyl-8-hydroxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 6-(2,3-Difluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(1-Ethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile; [6-(4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)pyridin-2-yl]acetonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-phenylacetonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-4-fluoro-phenylacetonitrile; 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluorophenylacetonitrile; 2-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)phenylacetonitrile; 6-(3-

AHPWA1DUSA

Fluoro-4-methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile; 6-(6-Amino-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazine-2-one; 6-(5-Diethoxymethyl-furan-3-yl) -4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; and 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbaldehyde; or a pharmaceutically acceptable salt thereof.

48(Original). A method of hormone replacement therapy, the method comprising administering to a mammal in need thereof a compound of the formula (I):



I

wherein:


R¹ and R² are independent substituents selected from the group consisting of H and C₁ to C₆ alkyl;

R³ is H;

R⁴ is H;

R⁵ is selected from the group consisting of (i) and (ii):

(i) a substituted benzene ring having the substituents X, Y and Z as shown below:



wherein:

AHPWA1DUSA

X is selected from the group consisting of halogen, CN, C₁ to C₃ alkyl, and substituted C₁ to C₃ alkyl;

Y and Z are independent substituents selected from the group consisting of H, halogen, CN, NO₂, and C₁ to C₃ alkyl; and

(ii) a five or six membered ring having in its backbone 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and containing one or two independent substituents selected from the group consisting of H, halogen, and CN;

wherein when R⁵ is a five-membered ring having in its backbone a NR⁶ heteroatom, and when R⁵ is attached at the two position of said ring, there is no CN substituent in the five position on said ring;
or pharmaceutically acceptable salt thereof.

49(Original). A compound selected from the group consisting of 6-(3-Methoxyphenyl)spiro[4H-3,1-benzoxazine-4,1-cyclobutan]-2(1H)-one; 8-Bromo-6-(3-chloro-4-fluorophenyl)-4,4-dimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 3-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 5-(8-Bromo-4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-2-fluorobenzonitrile; 6-(3-Bromophenyl)-1,4,4-trimethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 6-(3-Fluorophenyl)-4-methyl-1,4-dihydro-2H-3,1-benzoxazin-2-one; 3-(4,4-Dimethyl-8-methoxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 3-(4,4-Dimethyl-8-hydroxy-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-5-fluorobenzonitrile; 6-(2,3-Difluoro-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(1-Ethyl-4,4-dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-5-fluoro-benzonitrile; [6-(4,4-dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)pyridin-2-yl]acetonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-5-fluoro-phenylacetonitrile; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-4-fluoro-phenylacetonitrile; 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-2-fluorophenylacetonitrile; 2-(4,4-

AHPWA1DUSA

Dimethyl 2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile; 6-(3-Fluoro-4-methoxy-phenyl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)phenylacetonitrile; 6-(6-Amino-pyridin-3-yl)-4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazine-2-one; 6-(5-Diethoxymethyl-furan-3-yl) -4,4-dimethyl-1,4-dihydro-benzo[d][1,3]oxazin-2-one; 4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]oxazin-6-yl)-furan-2-carbaldehyde; N-[4-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-3,1-benzoxazin-6-yl)-2-fluoro-phenyl]-acetamide; 3-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)benzenesulfonamide; 5-(4,4-Dimethyl-2-oxo-1,4-dihydro-2H-benzo[d][1,3]-oxazin-6-yl)-thiophene-2-sulfonamide; and 4-(1,4-Dihydro-4,4-dimethyl-2-oxo-2H-3,1-benzoxazin-6-yl)-2-furancarboxaldehyde oxime;
or a pharmaceutically acceptable salt thereof.